



**NISSEN FUNDOPLICATION
DISCHARGE INSTRUCTIONS**

PLEASE REVIEW THESE INSTRUCTIONS THOROUGHLY BEFORE SURGERY

PREPARING FOR SURGERY

You will receive a call from the Preadmit/Same Day Surgery Nurse to review your health status, discuss any medications you are currently taking, and receive instructions about your surgery. These instructions are critical to your well-being. Please follow these guidelines completely or your surgery may have to be delayed or canceled.

Preadmit/Same Day Surgery Nurse: _____

Phone Number of Preadmit/Same Day Surgery Nurse for Questions: _____

Surgery Date/Time: _____

Arrival Time for Surgery:* _____

*Arrive for surgery at our Maine Street entrance. Check in at the Registration desk.

Medications **to take** the morning of Surgery: _____

Medication **to stop** prior to or the morning of Surgery: _____

Notes for Surgery: _____

SPECIAL INTRUCTIONS FOR YOUR SURGERY

Stop taking Aspirin **unless** otherwise directed.

Your Surgeon has scheduled a Pre-Surgery (Pre-Op) Appointment on _____.

Please arrive at Admissions, Arkansas Street entrance, to register. Volunteers can provide directions and/or assistance as needed to guide you to the appropriate departments for your tests.

Other special instructions:

Please understand that your surgery time may change due to emergency circumstances. A Preadmit/Same Day Surgery Nurse will call you 24 hours before your scheduled surgery date to verify both your arrival and surgery time.

Make arrangements for someone to remain present during surgery and to drive you home after your procedure. It is best to have a caregiver/family member remain with you for the first twenty-four hours after your surgery.



DAY OF SURGERY

Before you leave home

Do not eat any food, drink any beverage (including ice, etc.), chew gum or use cough drops/throat lozenges, etc. after 12:00 a.m./Midnight the day of surgery, unless directed. You may brush your teeth or rinse your mouth with a small amount of water; however, do not swallow.

Remove any make-up or nail polish. If you wear contact lenses, remove them and use your eyeglasses. Wear loose, comfortable clothing and shoes.

All jewelry should be removed. It is best to leave all valuables and jewelry at home or with your caregiver/family member.

Bring your photo identification and health insurance card(s).

Arriving at LMH

Check in at the Registration desk just inside the Maine Street entrance at LMH. The registration and nursing staff will prepare you for your procedure. You will have the opportunity to speak with anesthesia and your surgeon before surgery, and we will keep your caregiver/driver up-to-date on your health while you are in surgery.

AFTER SURGERY

Depending on your procedure, you will be released to home from the Recovery Unit, or be transferred to the Post-Surgical Care Floor (Third Floor, West). Once you are home, a nurse will contact you for a health update and to review your discharge instructions. This information is also available through My Patient Portal. You can access this site through our main web page at www.lmh.org/my-patient-portal/.

If you require FMLA paperwork, this may be returned to you via portal too. Please allow a week for completion.

Call the Lawrence General Surgery office at (785) 505-2200 at your first opportunity to schedule a follow-up appointment for 7-14 days after your surgery.

Follow-Up Surgery Appointment
with Lawrence General Surgery: _____

Call at any time if you have questions or notice any signs of infection, experience a fever over 101 degrees, wound redness, increasing pain or swelling, or an odorous, thick discharge. In case of emergency, please call 911 or go to the nearest emergency department.



Call at any time if you have questions or notice any signs of infection, experience a fever over 101 degrees, wound redness, increasing pain or swelling, or odorous, thick discharge. In case of emergency, please call 911 or go to the nearest emergency department.

OPERATIVE SITE

There may be swelling, bruising and a small amount of drainage at the operative area(s). Apply an ice pack to the operative area(s) intermittently when you get home for the next 1-2 days. The ice pack should not be applied directly to the skin (place a cloth between the ice pack and the area). The ice pack should not be left in place more than 15-30 minutes at a time.

You may have a tender lump at each incision site. This will resolve during normal healing over the next 4-6 weeks.

Abdominal fullness and discomfort into the back and shoulder area are common for a few days. Vomiting or belching is difficult if not impossible after this procedure. Talk with your doctor if you have questions.

Skin Adhesive Dressing

No bandage is needed unless the incision starts to drain or ooze. Do not use ointment on the skin adhesive. You may shower the same day as your surgery, but do not soak the incision area(s) for 1-2 weeks.

Steristrip™ Dressing

DO NOT remove the Steristrip™ tape(s) from the incision. Keep the area dry for 24 hours. You may shower after this time, but do not soak the incision area(s) for 1-2 weeks. Please keep the dressing in place for 24-48 hours. If you experience draining or the tape causes a skin irritation, please contact the office.

Use additional dressing (gauze, Band-Aid®, etc) as needed to cover any drainage from the incision.

PAIN CONTROL

After the local anesthetic wears off, pain medication may be needed. Begin taking pain medication as soon as discomfort begins and as directed thereafter. Pain medications may have side effects like nausea, vomiting and constipation. Please call the office if you have questions about your pain control.

Mild Pain

Acetaminophen (Tylenol®) ibuprofen (Advil® or Motrin®) or naproxen (Aleve®) are acceptable medications to take for mild pain.

Severe Pain

Take the prescription medication provided by your doctor in addition to a form of ibuprofen (Advil® or Motrin®).

DO NOT drive or drink alcohol while taking prescription pain medication.

DO NOT take ibuprofen (Advil® or Motrin®) or naproxen (Aleve®), if it bothers your stomach, you have kidney disease or you are taking Toradol (ketorolac).

DO NOT take acetaminophen (Tylenol®) while using other medications which also contain acetaminophen, such as Lortab or Percocet.

BOWEL and BLADDER FUNCTION

If you are unable to urinate within 6-8 hours of arriving at home, call your doctor. A stool softener (Colace® or equivalent) is often recommended for a few weeks after surgery. Laxatives (Milk of Magnesia or Miralax®) are permissible and often needed if you are taking prescription pain medication.



ACTIVITY

Avoid strenuous activity for the next four weeks. Do not lift more than 30 pounds for the next four weeks. Avoid repetitive bending, stooping and straining. Do not push or pull any item for the next four weeks.

Light activities such as walking, desk work, climbing stairs, or driving can be resumed as tolerated.

Sexual intercourse is typically permissible based on your comfort level or after your post operative.

You may return to work as soon as you can perform your job activities without causing pain. Typically this is light duty work after the first week, and manual labor in one month.

DO NOT drive if prescription pain medication is being used.

DIET

You will resume your normal diet three weeks after surgery. Until then, follow a mechanical diet which provides foods that are easier to swallow. It is important to take small bites and chew foods thoroughly before swallowing. (A Mechanical Diet Guide will be provided to you.) Also increase your fluid intake. Drink an extra 5-6 glass a water each day.

PATIENT PORTAL

These discharge instructions, plus other valuable information – visit summary, certain test results, FMLA or Return to Work forms, etc. – are available through My Patient Portal. You can access this site through our main web page at www.lmh.org/my-patient-portal/. You can also request appointments and ask your doctor questions through this secure, anytime access to personal health information.